


AD-A261 089



PY

KEEP THIS COPY FOR REPRODUCTION PURPOSES

2

## MENTATION PAGE

Form Approved

OMB No. 0704-0188

estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, S.W., Washington, D.C. 20540, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 11/12/92	3. REPORT TYPE AND DATES COVERED Final Report 3/15/89 - 6/14/92	
4. TITLE AND SUBTITLE EFFECT OF CYTOSKELETAL REAGENTS ON STRETCH ACTIVATED ION CHANNELS		5. FUNDING NUMBERS DAAL03-89-k-0064	
6. AUTHOR(S) Dr. Frederick Sachs		7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biophysical Sciences Dept. 120 Cary Hall SUNY at Buffalo Buffalo, NY 14214	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709-2211		8. PERFORMING ORGANIZATION REPORT NUMBER ARO 26099.9-15	
11. SUPPLEMENTARY NOTES The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			
13. ABSTRACT (Maximum 200 words) Mechanically sensitive ion channels have been proposed to respond to membrane tension. We have developed methods to measure the membrane tension in a patch and have examine the stress strain relationship. The results show that, in general, tension is the critical variable. However, we have also found one channel in glial cells whose gating is additionally dependent upon the curvature of the membrane. Stresses in a patch caused by suction lead to lipid flow along the wall of the pipette, but the flow is constrained by the highly extensible cytoskeleton whose area elastic constant is ca. 50dyn/cm.  We developed tools to use high voltage electron microscopy to study the structure of patches and have characterized the placement of cytoskeleton, lipids and receptors in a variety of preparations. We also developed a new algorithm to align projections for tomography in order to examine the three dimensional structure of patches.  We have demonstrated mechanically induced release of calcium via stretch activated ion channels in heart cells extending work on the molecular level to the whole cell level.			
14. SUBJECT TERMS mechanical transduction, patch, calcium, electron microscopy light microscopy, tomography		15. NUMBER OF PAGES 5	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
		20. LIMITATION OF ABSTRACT UL	

93 2 1 0 5 5

## Effect of Cytoskeletal Reagents on Stretch Activated Ion Channels

Dr. Frederick Sachs

November 6, 1992

US. Army Research Office  
DAALO3-89-0064

SUNY  
Buffalo, NY 14214

Approved for Public Release

**DTIC QUALITY INSPECTED 3**

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input checked="checked" type="checkbox"/>
Unannounced	<input checked="checked" type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

## Problem Studied:

What makes cells mechanically sensitive? This work examined the mechanical properties of patch clamped cell membranes with regard to both structure, stress-strain relationships and properties of the mechano-electric transduction channels.

## Major results:

We have found that patch clamped membranes always are formed with the cytoskeleton attached. this has been verified by 2D and 3D electron microscopy. We have developed and published procedures for making such images. Working with light microscopy, we have found that the cytoskeletal structure of patches is highly variable. Sometimes the cytoskeleton sticks in a compact layer to the membrane and there is little Brownian motion of entrained particles. In other cases the cytoplasm is disrupted and particles diffuse readily. The variability of the structures accounts, at least in part, for the variability of stretch induced responses in cells.

By combining electrophysiology with light microscopy of patches we have been able to measure the elasticity of membranes and to clearly separate the properties of lipids from those of the cytoskeleton. In skeletal muscle cells the cytoskeleton has an area elastic constant of about 50dyn/cm and this value is insensitive to treatment with cytochalasin suggesting that actin is not tension bearing in this membrane. In heart cells, on the other hand, the elasticity decreases to 25dyn/cm with cytochalasin. We were able to show that under stress, the lipids in a membrane patch are free to flow along the walls of the pipette. We were able to show that a major source of error in the measurement of stretch activated channel sensitivity can be removed by using as the independent variable tension rather than applied pressure.

We studied mechanical transduction in a variety of cells types and found mechanically sensitive channel in the lateral walls of auditory hair cells from the guinea pig. These channels are not the normal transducers responsible for the major sensory activity but may play a role in tuning. These channels may explain why hair cells will contract when a frequency modulated water jet is applied to the cell body.

We found in glial cells that there are multiple types of transduction channels and one of them has the unique property of only responding when the membrane is curved toward the cytoplasm. This is the first observation of such a curvature sensitive channel.

On a cellular level we have shown that mechanical stimulation of heart cells can cause the release of intracellular calcium. The mechanism of this coupling probably involves stretch activated ion channels since the response is blocked by low extracellular calcium, blockage of the channels with Gd ions or suppression of channel expression by omission of growth factors from the culture medium.

## Participating Scientific Personnel:

Dr. Frederick Sachs, Dr. Charles Bowman, Dr. Masahiro Sokabe, Dr. Wade Sigurdson, Dr. Zhonqi Jing, Dr. R.J. Salvi, Dr. J.P. Ding, Dr. A. Ruknudin, Dr. X.C. Yang (obtained his PhD during the project),

## Publications:

1. DING, J. P., C. L. BOWMAN, M. SOKABE, AND F. SACHS. Mechanical transduction in glial cells: SICS and SACS. *Biophys. J.* 55: 244a, 1989.

2. GAO, X. AND F. SACHS. Improving performance of motorized slides for micromanipulation. *J. Neurosci. Methods* 28: 225-227, 1989.
3. RUKNUDIN, A., M. SONG, A. AUERBACH, AND F. SACHS. The structure of patch clamped membranes in high voltage electron microscopy. *Proc. Elec. Mic. Soc. Am.* 47: 936-937, 1989.
4. SACHS, F. Stretch-activated ion channels as acoustical transducers. *J. Acoust. Soc. Am.* 85: S29, 1989.
5. SACHS, F. AND M. SOKABE. Stress and strain in patch clamped membranes. *Int. Conf. Video. Microscopy* 37, 1989.(Abstract)
6. SACHS, F. Ion Channels as Mechanical Transducers. In: *Cell Shape: Determinants, Regulation and Regulatory Role*, edited by W. D. Stein and F. Bronner. San Diego, NY, Berkeley, Boston: Academic Press, 1989, p. 63-92.
7. YANG, X-C. AND F. SACHS. Block of Stretch-Activated Ion Channels in *Xenopus* Oocytes by Gadolinium and Calcium Ions. *Science* 243: 1068-1071, 1989.
8. JING, Z. AND F. SACHS. *TACIT: a command line library for two and three dimensional image processing*. Copyright, SUNYAB. 1990, (UnPub)
9. MCEWEN, B. F., M. J. SONG, A. RUKNUDIN, D. P. BARNARD, J. FRANK, AND F. SACHS. Tomographic three dimensional reconstruction of patch clamped membranes imaged with the high voltage electron microscope. *XII Int. Conf. Electron Microscopy* 522-523, 1990.(Abstract)
10. SACHS, F. Stretch-sensitive ion channels. *Sem. Neurosci.* 2: 49-57, 1990.
11. SACHS, F. AND M. SOKABE. Stretch-activated ion channels and membrane mechanics. *Neurosci. Res. Suppl.* 12: S1-S4, 1990.
12. SOKABE, M. AND F. SACHS. The structure and dynamics of patch-clamped membranes: a study by differential interference microscopy. *J. Cell Biol.* 111: 599-606, 1990.
13. SOKABE, M. AND F. SACHS. Stress and strain in patch clamped membranes. *Biophys. J.* 57: 265a, 1990.
14. SOKABE, M. AND F. SACHS. Quantitative study of stress, capacitance, and sa-channel activity in patch clamped membranes. *Intl. Biophy. Cong.* 1990.(Abstract)
15. YANG, X. -C. AND F. SACHS. Characterization of stretch-activated ion channels in *Xenopus* oocytes. *J. Physiol. (Lond.)* 431: 103-122, 1990.
16. BUSTAMANTE, J. O., A. RUKNUDIN, AND F. SACHS. Stretch-activated channels in heart cells: Relevance to cardiac hypertrophy. *J. Cardiovasc. Pharmacol.* 17 Suppl. 2: S110-S113, 1991.
17. DING, J. P., R. J. SALVI, AND F. SACHS. Stretch-activated ion channels in guinea pig outer hair cells. *Hearing Res.* 56: 19-28, 1991.

18. IZU, Y. C. AND F. SACHS. B-D-Xyloside Treatment Improves Patch Clamp Seal Formation. *Pflugers Arch* 419: 218-220, 1991.
19. JING, Z. AND F. SACHS. Alignment of Tomographic Projections Using an Incomplete Set of Fiducial Markers. *Ultramicroscopy* 35: 37-43, 1991.
20. RUKNUDIN, A., M. J. SONG, AND F. SACHS. The ultrastructure of patch-clamped membranes: a study using high voltage electron microscopy. *J. Cell Biol.* 112: 125-134, 1991.
21. RUKNUDIN, A., J. O. BUSTAMANTE, AND F. SACHS. Stretch activated channels of chick cardiac myocytes. *J. Cell. Bioch. Suppl.* 15C: 197, 1991.(Abstract)
22. SACHS, F. AND H. LECAR. Stochastic models for mechanical transduction. *Biophys J* 59: 1143-1145, 1991.
23. SACHS, F., W. SIGURDSON, A. RUKNUDIN, AND C. BOWMAN. Single-channel mechanosensitive currents. *Science* 253: 800-801, 1991.
24. SACHS, F. Mechanical transduction by membrane ion channels: a mini review. *Mol. Cell. Biochem.* 104: 57-60, 1991.
25. SIGURDSON, W. AND F. SACHS. Mechanical stimulation of cardiac myocytes increases intracellular calcium. *Biophys J* 59: 469a, 1991.(Abstract)
26. SOKABE, M., F. SACHS, AND Z. JING. Quantitative video microscopy of patch clamped membranes - stress, strain, capacitance and stretch channel activation. *Biophys. J.* 59: 722-728, 1991.
27. BOWMAN, C. L., J. P. DING, F. SACHS, AND M. SOKABE. Mechanotransducing ion channels in astrocytes. *Brain Res.* 584: 272-286, 1992.
28. RUKNUDIN, A., F. SACHS, AND O. BUSTAMANTE. Stretch activated ion channels in chick heart cells. *Am. J. Physiol.* 1992.(In Press)
29. SACHS, F. Stretch sensitive ion channels: an update. In: *Sensory Transduction*, edited by D. P. Corey and S. D. Roper. NY: Rockefeller Univ. Press, Soc. Gen. Physiol., 1992, p. 241-260.
30. SIGURDSON, W. S., A. RUKNUDIN, AND F. SACHS. Calcium imaging of mechanically induced fluxes in tissue-cultured chick heart: role of stretch-activated ion channels. *Am. J. Physiol.* 262: H1110-H1115, 1992.
31. SOKABE, M. AND F. SACHS. Towards a molecular mechanism of activation in mechanosensitive ion channels. In: *Advances in Comparative and Environmental Physiology*, v10, edited by F. Ito. Berlin: Springer-Verlag, 1992, p. 55-77.

Report of Inventions: None